

EMBEDDING VERSUS IMMERSION IN GENERAL RELATIVITY

EDMUNDO M. MONTE*

Departamento de Física, Universidade
Federal da Paraíba, 58059-970, João Pessoa, Paraíba, Brazil.

August 23, 2009

Abstract

We briefly discuss the concepts of immersion and embedding of space-times in higher-dimensional spaces. We revisit the classical work by Kasner in which he constructs a model of immersion of the Schwarzschild exterior solution into a six-dimensional pseudo-Euclidean manifold. We show that, from a physical point of view, this model is not entirely satisfactory since the causal structure of the immersed space-time is not preserved by the immersion.

1 Introduction

The first example of immersion of space-times in higher-dimensional spaces came from Kasner, in 1921, when he explicitly exhibited a local isometric immersion of the Schwarzschild exterior solution.[1] Interest in higher-dimensional theories of space-time seems to be increasing these days. Actually, the ideas that underlie the modern developments were first put forward by Kaluza and Klein, who considered the possibility that our spacetime may have extra dimensions.[2] Currently there is a huge literature on the subject of extra-dimensional theories, immersion and embedding of space-times, mainly motivated by the generalized Kaluza-Klein theory, string theory and the recent braneworld scenario.[3, 4, 5, 6, 7] In all these theoretical developments the geometrical concepts of immersion and embedding play a very important role.[8] Our main goal in this paper is to give a precise meaning

*e-mail: edmundomonte@pq.cnpq.br and edmundomonte@pq.cnpq.br

of these concepts and briefly discuss the immersion obtained by Kasner. We prove that in this case the immersion does not preserve the causal character of curves, thereby not being quite appropriate as a physical model.

2 Immersions and Embeddings

Let (V_N, g) and (V_D, η) be two differentiable manifolds V_N and V_D , with dimensions N and D , endowed with the metric tensors g and η , respectively. A differentiable application $Y : (V_N, g) \rightarrow (V_D, \eta)$ is called a *local and isometric immersion* if:

$\forall p \in U \subset V_N$, $dY_p : T_p V_N \rightarrow T_{Y(p)} V_D$ is injective, where U is an open neighbourhood of p , and $g(v, w) = \eta(dY_p(v), dY_p(w))$, $\forall v, w \in T_p V_N$. If, in addition, Y is a homeomorphism onto $Y(U)$, where $Y(U)$ has the induced topology of V_D , then Y is called a *local and isometric embedding*. When $V_N \subset V_D$ and the inclusion $Y : V_N \subset V_D \rightarrow V_D$ is an embedding, we say that V_N is a *submanifold* of V_D . [9]

Let us now consider (V_4, g) as a pseudo-Riemannian manifold, corresponding to a solution of Einstein's equations, locally and isometrically embedded in another pseudo-Riemannian manifold (V_D, η) . In local coordinates $\{x^i\}$ and $\{Y^\mu\}$ of V_N , V_D , the isometric condition takes the form

$$g_{ij} = \eta_{\mu\nu} Y_{,i}^\mu Y_{,j}^\nu, \quad (1)$$

where g_{ij} is the induced metric on V_4 ; $Y_{,i}^\mu = \frac{\partial Y^\mu}{\partial x^i}$ and the indices now take value in the range $\mu = 1, \dots, D$; $i = 1, \dots, 4$.

From the above definitions we see that the essential distinction between immersion and embedding is of topological character. In fact, any embedding is an immersion, while the converse is not always true. [10]

3 The Immersion of Schwarzschild Space-time

The Schwarzschild exterior solution, which describes the space-time geometry outside a spherical symmetric body, is given, in spherical coordinates (t, r, θ, ϕ) by

$$ds^2 = (1 - 2m/r)dt^2 - (1 - 2m/r)^{-1}dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (2)$$

where m is the so-called geometric mass. Actually, one can regard the above geometry as defining two different space-times: [11, 12]

a) The exterior Schwarzschild space-time (V_4, g) , where $V_4 = P_I^2 \times S^2$;
 $P_I^2 = \{(t, r) \in \mathbb{R}^2 \mid r > 2m\}$, and
b) The Schwarzschild black hole (B_4, g) , with
 $B_4 = P_{II}^2 \times S^2$, $P_{II}^2 = \{(t, r) \in \mathbb{R}^2 \mid 0 < r < 2m\}$.
In both cases, S^2 denotes the sphere of radius r and metric induced from (2).

Now define the space $(E, g) = ([P_I^2 \cup P_{II}^2] \times S^2, g)$ and consider an application $Y : (V_4, g) \rightarrow (V_6, \eta)$, where the pseudo-Euclidean metric η is given by the line element: [1, 10]

$$ds^2 = dY_1^2 + dY_2^2 - dY_3^2 - dY_4^2 - dY_5^2 - dY_6^2. \quad (3)$$

It is not difficult to see that the application Y defined by the equations

$$\begin{cases} Y_1 = (1 - 1/r)^{1/2} \cos t \\ Y_2 = (1 - 1/r)^{1/2} \sin t \\ Y_3 = f(r), \quad (df/dr)^2 = \frac{1+4r^3}{4r^3(r-1)} \\ Y_4 = r \sin \theta \sin \phi \\ Y_5 = r \sin \theta \cos \phi \\ Y_6 = r \cos \theta, \end{cases}$$

represents a local isometric immersion of (V_4, g) into (V_6, η) . For convenience we are taking $2m = 1$ in the Schwarzschild metric. Indeed, a simple calculation shows that (1) holds and that $dY_p : T_p V_4 \rightarrow T_{Y(p)} V_6$ is injective, i.e.,

$$\dim[dY_p(T_p(V_4))] = 4,$$

or, equivalently,

$$\det(\text{minor of matrix}[dY_p]_{6 \times 4}, 4 \text{ order}) = m \sin^2 \theta \sin \phi \neq 0,$$

for $0 < \theta < \pi$ and $0 < \phi < \pi$ or $\pi < \phi < 2\pi$.

It is easy to see that this isometric immersion does not constitute an isometric embedding. Indeed, to verify this fact just consider a curve

$$\begin{aligned} \alpha & : (a, b) \subset \mathbb{R} \longrightarrow (V_4, g) \\ t & \rightarrow \alpha(t) \in (U \subset V_4, g) \end{aligned}$$

The image of $\alpha(t)$ through the application Y is $Y(\alpha(t)) = (Y_1(\alpha(t)), \dots, Y_6(\alpha(t)))$. Now, if we take $a < t_1 < t_2 < b$, since Y_1 and Y_2 are periodic functions of t ,

it is clear that it is possible to have

$$\alpha(t_1) \neq \alpha(t_2) \text{ and } Y_\mu(\alpha(t_1)) = Y_\mu(\alpha(t_2)), \mu = 1, 2. \quad (4)$$

In other words, there is a neighbourhood $U \subset V_4$, where Y is not a homeomorphism. In the particular case where the curve $\alpha(t)$ is timelike it follows directly from (4) and the isometric condition (1) that $Y(\alpha(t))$ is a closed timelike curve in $Y(U) \subset (V_6, \eta)$. Obviously, from the standpoint of physics this state of affairs is not entirely satisfactory as the causal structure of the space-time (V_4, g) is not preserved by the immersion.

It is natural at this point to wonder whether an isometric embedding of the Schwarzschild space-time into a six-dimensional has ever been found. The positive answer to this question was given in 1959, with the work of Fronsdal, who found a regular embedding of the Schwarzschild exterior solution into a six-dimensional pseudo-Euclidean space by using hyperbolic functions instead of the trigonometric functions used by Kasner.[13]

4 Comment

Finally, we conclude that, even though any immersion can be considered locally as an embedding, it is important to be aware of the existence sometimes of pathological domains in the immersed space-time, as in the case of Kasner's immersion of the Schwarzschild exterior solution. Therefore one should be careful with the terminology. From the point of view of physics, models that employs embeddings are much useful than those that employs immersions, since the latter do not preserve the topology of the immersed manifold.

Acknowledgments

We would like to thank Professor Marcos Maia for useful discussions. Thanks also go to FAPES-ES/CNPq (PRONEX) and FAPESQ-PB/CNPq (PRONEX) for financial support. We are indebted to the referees, whose comments helped us to improve the manuscript.

References

- [1] E. Kasner, *Am. J. Math.* **43**, 130 (1921).

- [2] T. Kaluza, *Sitzunger. Preuss. Akad. Wiss. Berlin, Phys. Math.* **K1**, 33 (1921). O. Klein, *Z. Phys.* **37**, 895 (1926).
- [3] Nima Arkani-Hamed *et al.*, *Phys. Lett. B* **429**, 263 (1998).
- [4] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 3370 (1999).
- [5] K. Maeda *et al.*, *Phys. Rev. D* **62**, 24012 (2000).
- [6] M. D. Maia and E. M. Monte, *Phys. Lett. A* **297**, 9 (2002).
- [7] R. Kerner, *Ann. Inst. H. Poincaré* **9**, 141 (1968).
- [8] For a nice review on embeddings and immersions, see, for instance, H. Goenner, *Local Isometric Embedding of Riemannian Manifolds and Einstein's Theory of Gravitation*, in *General Relativity and Gravitation*, **I**, (A. Held, Plenum Press, New York, 1980). See also, J. Rosen, *Rev. Mod. Phys.*, **37**, 204 (1965).
- [9] M. P. Carmo, *Riemannian Geometry*, (Birkhuser, Boston, 1993).
- [10] M. D. Maia and E. M. Monte, *Mat. Contemp.*, **13**, 229 (1997).
- [11] S. Hawking and G. Ellis, *The Large Scale Structure of Space-Time*, (Cambridge Univ. Press, Cambridge, 1973).
- [12] B. O'Neill, *Semi-Riemannian Geometry*, (Academic Press, New York, 1970).
- [13] C. Fronsdal, *Phys. Rev.*, **116**, 778 (1959).